

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Exteris Stressgard®

**TURF FUNGICIDE**

**ACTIVE CONSTITUENTS:** 12.5 g/L FLUOPYRAM  
12.5 g/L TRIFLOXYSTROBIN

**GROUP 7 11 FUNGICIDE**

**5 L** For control of various diseases in turf as specified in the Directions for Use.

 **CENTAUR**  
ASIA PACIFIC  
[www.centaur-asiapacific.com](http://www.centaur-asiapacific.com)  
[info@centaur-asiapacific.com](mailto:info@centaur-asiapacific.com)

**DIRECTIONS FOR USE****RESTRAINTS**

DO NOT apply by aircraft

**SPRAY DRIFT RESTRAINTS**

**DO NOT** apply with spray droplets smaller than a **MEDIUM** spray droplet size category according to nozzle manufacturer specifications that refer to ASAE S572 Standard to the BCPC Guideline.

**DO NOT** apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

**DO NOT** apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are:

1. date and start and finish times of application;
2. location address and areas sprayed;
3. full name of this product;
4. amount used per hectare and number of hectares applied to;
5. situation and disease;
6. wind speed and direction during application;
7. air temperature;
8. nozzle brand, model and type and spray system pressure measured during application;
9. name and address of person applying this product.

(Additional record details may be required by the State or Territory where this product is used).

**MANDATORY NO-SPRAY ZONES**

**DO NOT** apply if there are aquatic and wetland areas, including aquacultural ponds or surface streams and rivers, within 5 metres downwind from the application area.

CROP	DISEASE	RATE	CRITICAL COMMENTS
Turf	Fusarium or Microdochium patch ( <i>Microdochium nivale</i> );  Dollar spot ( <i>Sclerotinia homoeocarpa</i> );  Helminthosporium disease ( <i>Bipolaris</i> spp., <i>Drechslera</i> spp., <i>Exserohilum</i> spp.);  Brown patch ( <i>Rhizoctonia solani</i> );  Gray leaf spot ( <i>Pyricularia grisea</i> );  Rust ( <i>Puccinia</i> spp.)	10 L/ha  Water volume: 400-800 L/ha	Apply on a 14-28 day spray interval to established turf when conditions are favourable for disease development. Use shorter spray interval under high disease pressure or for early curative treatment.  For Brown Patch control, apply sufficient irrigation to wash the product into the crown and upper root zone.  Apply a maximum of four applications per year.  Apply a maximum of 40 L / ha per year.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD**

DO NOT graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock.

**GENERAL INSTRUCTIONS****Application**

Good control requires good coverage. Application should be made using sufficient water to ensure thorough coverage.

**IPM Compatibility**

Risk to non-target insects: Exteris may have adverse effects on some non-target beneficial insects, in particular predatory bugs (e.g. *Orius* spp.). Safety to ladybirds and syrphid flies has not been evaluated. This risk should be taken into account where integrated pest management utilising these organisms is practised.

**Compatibility**

As formulations of other manufacturers' products are beyond the control of Bayer, all mixtures should be tested prior to mixing commercial quantities.

**RESISTANCE WARNING**

<b>GROUP</b>	<b>7</b>	<b>11</b>	<b>FUNGICIDE</b>
--------------	----------	-----------	------------------

Exteris Stressgard Turf Fungicide is a combination of a succinate dehydrogenase inhibitor (SDHI) and a quinone outside inhibitor (QoI) fungicide. For fungicide resistance management Exteris Stressgard is both a Group 7 and a Group 11 fungicide. Some naturally occurring individual fungi resistant to Exteris Stressgard and other Group 7 and Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Exteris Stressgard and other Group 7 and Group 11 fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of Exteris Stressgard to control resistant fungi.

**PRECAUTIONS**

**Re-entry period** Do not allow entry into treated areas until the spray has dried.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Very toxic to aquatic life. DO NOT contaminate wetlands or water courses with this product or used containers.

**PROTECTION OF LIVESTOCK**

DO NOT graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

**SAFETY DIRECTIONS**

May irritate the eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes. When handling product, wear chemical resistant gloves. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26)

**SAFETY DATA SHEET**

Additional Information is listed in the Safety Data Sheet which is available at [www.environmentalscience.bayer.com.au](http://www.environmentalscience.bayer.com.au).

**Exclusion of Liability**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No.: 83704 / 108864

Exteris Stressgard® is a Registered Trademark of the Bayer Group

Bayer CropScience Pty Ltd  
A.B.N. 87 000 226 022  
Level 1, 8 Redfern Rd  
Hawthorn East, Vic. 3123  
Technical enquiries: 1800 804 479  
Website: [www.environmentalscience.bayer.com.au](http://www.environmentalscience.bayer.com.au)

FOR 24 HOUR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 033 111