

EverGreen 18-5-0 Minis + 5% Iron with 0.40% Barricade®

For Preemergence Control of Weed Grasses and Broadleaf Weeds in all Established Turf Grasses, Lawns, Golf Course Fairways, and Sod Farms, (not for putting greens). Use for Preemergence Weed Control Around Landscape Ornamentals Including Established Perennials and Wildflower Plantings.

*CAS (2909-21-2) EPA Reg. No. 7001-385 Barricade® is a Reg. Trademark of Syngenta.

EPA Est. No. 61842-CA-1

KEEP OUT OF REACH OF CHILDREN CAUTION

 GUARANTEED ANALYSIS: 18-5-0

 Total Nitrogen (N)
 18.00%

 18.00% Ammoniacal Nitrogen
 5.00%

 Available Phosphate(P₂O₅)
 5.00%

 Iron (Fe)
 18.00%

 Derived from Ammonium Sulfate, Ammonium Phosphate, and Iron Oxysulfate.
 5.00%

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS and DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reaction in some individuals.

FIRST AID					
IF SWALLOWED:	Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.				
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. 				
IF ON SKIN OR CLOTHING:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. 				
IF INHALED:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. 				

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment advice.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicator and other handlers must wear long-sleeved shirt and long pants; chemical resistant gloves; shoes plus socks; a dust mask and protective eyewear. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

<u>Users should:</u> Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

READ ENTIRE LABEL BEFORE USING

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on the farms, forests, and nurseries and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (sod farms, commercial nurseries).

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves, such as butyl rubber > 14 mils or natural rubber > 14 mils or Nitrile rubber > 14 mils.
- Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The "WPS" applies when this product is used to produce agricultural plants on farms, forests, or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, chemical resistant gloves, shoes with socks, and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until dusts have settled and the turf grass or soil is dry.

CRABGRASS SEED GERMINATION DATES						
Approximate Dates						
1. Mid-north & northwestern	After May 30					
Upper mid-states & northeast	After May 10					
Lower mid-states	After April 20					
Upper southern states	After Mar 20					
5. Warm season states	Jan 1 to Mar 20					

This product is a turf fertilizer and a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses, golf courses (excluding putting greens), for sod farms, and lawns, and around landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least ½ inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control.

APPLICATION DIRECTIONS

mly with suitable calibrated application equipment. Do not apply otherwise.

ESTÁBLISHEÐ TURF


Zoysia

This product is a preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds listed below in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the Maximum Annual Rates Chart of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least ½ inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates.

The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the weeds listed below. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of "residual preemergence weed control" provided by this product is related to the rate applied.

This product is recommended for use on the turf grass species listed in the following chart Do not exceed the maximum yearly rate as given in this chart. 1) Maximum application rate of this product per calendar year by turf grass species **Turf Species** Lbs. Product Lbs. Product Lbs. A.I./Acre / 1,000 Sq. ft. / Acre Creeping Bentgrass 162 lbs. 0.65 lb. Creeping Red Fescue 0.75 lb. 187 lbs. Buffalograss 250 lbs. 1.00 lb. Kentucky Bluegrass Perennial Ryegrass 375 lbs. 8.6 1.50 lbs. Bermudagrass (2) Bahiagrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf type)



- (1) These are the maximum rates per calendar year by species limitations.
- (2) May be used on newly-sprigged or plugged Bermudagrass at rates not to exceed 130 lbs./A. Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.
- •Do not apply more than 375 lbs. of this product to the same area per calendar year.

•Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Chart.

When used in accordance with this label, this product will provide control of the following weeds: Barnyardgrass, Bluegrass, Annual (Poa Annua) Carpetweed, Chickweed, Common Chickweed, Mouseear (from seed), Crabgrass (Large, Smooth), Crowfootgrass, Cupgrass Woolly, Foxtails, Annual, Goosegrass (1), Henbit, Itchgrass, Johnsongrass (from seed), Junglerice, Knotweed, Kochia, Lambsquarter, Common, Lovegrass, Panicum (Texas, Fall, Browntop); Pigweed; Purslane, Common; Pusley, Florida, Rescuegrass, (3), Shepherd's purse (2); Signalgrass, Broadleaf; Speedwell, Persian Sprangletop, Spurge, Prostrate; Witchgrass;

- (1) In many areas a single application of 162 to 375 lbs. per acre will control goosegrass. However under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 162 to 250 lbs. per acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rates Chart**. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rates Chart on this label.
- (2) Application for Shepherd's purse should be made in late summer, fall, or winter prior to germination.
 (3) Suppression only of rescuegrass. Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

 WHEN TO: APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil

WHEN TO: OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding and/or reseeding results.

Product Rate Lbs./Acre	Lbs. Product/ 1,000 Sq. ft.	Months Before Overseeding			
		A. I./Acre	North	Transition	South
125	2.87	0.50	4	4	4
162	3.72	0.65	5	4	4
187	4.29	0.75	6	5	5
200	4.59	0.80	-	6	6
250	5.74	1.00	-	7	7
287	6.59	1.15	-	-	9
325	7.46	1.30	-	-	10
375	8.60	1.50	-	-	12

Application Equipment and Spreader Settings

The size and bulk density of granules in this product may vary; therefore, calibrate your granular applicator prior to application of this product. Follow directions in the owner's manual for calibrating your spreader. The settings indicated in the following table should only be used as a guide in calibrating your spreader.

SPREADER SETTINGS

V. 112/15/21/11/10/								
1/2 lb	3/4 lb	1 lb						
2.87	4.29	5.74						
3 1/2 A	4 1/4 B	4 1/2 A						
3.5	3.9	4.2						
14	16	18						
11	13 1/4	15 1/4						
I	K 1/4	M						
4 1/2	5 1/4	6						
32	39	44						
	3 1/2 Ib 2.87 3 1/2 A 3.5 14 11 1 4 1/2	3/4 lb 2.87 4.29 3 1/2 A 4 1/4 B 3.5 3.9 14 16 11 13 1/4 I K 1/4 4 1/2 5 1/4						

LANDSCAPE ORNAMENTALS INCLUDING ESTABLISHED WILD FLOWER PLANTINGS

This product may be applied for residual preemergence weed control in Ornamentals as listed.

Apply at 162 to 375 lbs. per acre (3.72 lbs. to 8.61 lbs. per 1,000 sq. ft.) of landscape plants, in fall and/or spring. Use the higher rates of application for longer control periods. Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 375 lbs. per acre.

Apply uniformly to the surface of the soil in ornamental planting beds with a granular applicator. Calibrate your applicator to assure appropriate dosage as described above under "Use Rates", prior to application. Calibrate your applicator as described in the owner's manual. The bulk density of this product may vary; therefore, to assure that the product is applied at the rates described, you must use a calibrated applicator. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least ½ inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

This product may be applied to established ornamentals as a broadcast, over-the-top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil. **Newly Transplanted Landscape Ornamentals**

This product may be applied to newly transplanted ornamentals as a broadcast, over-the-top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. Irrigate after application as directed above.

This product will not harm most trees, shrubs, vines and flowers. The species listed here are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface. Avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over-the-top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

applying to manierial shoots which may delieal the effort to get uniform distribution on the soil surface. Another produce is uniformly applied to the soil surface. Another produces in uniform distribution on the soil surface. This product may be applied over-the-top of these species. When plants are under stress (such as heat, applied to the soil surface). Another products are under stress (such as heat, applied to the soil surface). Another products are under stress (such as heat, applied to the soil surface). Another products are under stress (such as heat, applied to the soil surface). Another products are under the surface and the surface an

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. If product is spilled sweep up and use as per label instructions or dispose of as indicated above.

Read the entire Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability,

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Injury, neffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of J.R. Simplot Company. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

Disclaimer of Warranties: To the extent consistent with applicable law, there are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of J.R. Simplot Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, J.R. Simplot Company disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

<u>LIMITATION OF WARRANTIES:</u> To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product,


whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at J.R. Simplot Company's option, the replacement of the product.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-jr.com

Simplot® and BEST® are registered trademarks of J.R. Simplot Company.

74347_evergreen1850minis+5%iron_0.4%barricade_EPA Notification 1-31-2011

NET WEIGHT 50 LB. BAG (22.68-Kg.)





Hong Kong | Singapore | Malaysia | Australia

Email: info@centaur-asiapacific.com

ASIA PACIFIC Website: www.centaur-asiapacific.com



MANUFACTURED IN U.SA. J. R. SIMPLOT COMPANY P.O. Box 198 • Lathrop, CA 95330