

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Recruit 200 SC

## INSECTICIDE

**ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID**

**GROUP 4A INSECTICIDE**

**For the control of various pests of certain vegetables and turf as specified in the Directions for Use table.**

**IMPORTANT: READ THIS LEAFLET BEFORE OPENING OR USING**



**Indigo Specialty Products Pty Ltd**  
ABN 15 631 459 660  
6/163-173 McEvoy Street,  
Alexandria NSW 2015  
Ph: 0402 725 887

Distributed by



**Contents**  
**1 L**

## DIRECTIONS FOR USE

### FOLIAR SPRAY APPLICATIONS

Crop	Insect	Rate	WHP	Critical Comments
Cucurbits	Green Peach Aphid	25 mL/100 L or 300 mL/ha	1 day	Apply at first sign of aphid infestation.
Capsicum Eggplant Potato			7 days	
Tomato			3 days	
Brassicac	Grey Cabbage Aphid, Turnip Aphid		7 days	Apply at first sign of aphid infestation. Add a wetting agent.
Sweet potato	Silverleaf Whitefly including type B	25 mL/100L or 250 mL/ha	7 days	Apply at first sign of whitefly or melon thrips infestation.
Cucumber			1 day	Apply dilute sprays (25 mL/100 L) to run off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Eggplant	Melon Thrips		7 days	
Turf	First instar larvae of: African Black Beetle, Argentinian Scarab, Pruinose Scarab	2.5 L/ha or 25 mL/100L Spray with at least 400 L water per hectare to ensure even coverage.  Preferably spray on to wet or dewy grass. Irrigate with 12 mm of water commencing within one hour of application.	-	Apply at peak egg hatch that is mid Spring to mid Summer depending on species.
	Larvae of Billbug			Monitor adult activity through late Spring and early Summer. Spray when numbers peak, or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS

**Cucurbits: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION**

**Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION**

**Brassicac, Capsicum, Eggplant, Potatoes, Sweet potatoes:**

**DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**

**DO NOT GRAZE ANY TREATED AREA, INCLUDING TURF, OR CUT FOR STOCK FOOD.**

**DO NOT FEED PRODUCE OR TURF CLIPPINGS FROM TREATED AREA TO ANIMALS, INCLUDING POULTRY.**

## GENERAL INSTRUCTIONS

### INSECTIDE RESISTANCE WARNING

For insect resistance management ProForce RECRUIT 200 SC Insecticide is a Group 4A Insecticide.

Some naturally occurring insect biotypes resistant to RECRUIT 200 SC and other group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if RECRUIT 200 SC or other Group 4A insecticides are used repeatedly. The effectiveness of RECRUIT 200 SC on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Indigo Specialty Products Pty Ltd accepts no liability for any losses that may result from the failure of RECRUIT 200 SC to control resistant insects. RECRUIT 200 SC may be subject to specific resistance management strategies. For further information contact your local supplier, Indigo Specialty Products Pty Ltd representative or local agricultural department agronomist.

**GROUP 4A INSECTICIDE**

### RESISTANCE MANAGEMENT STRATEGY

#### **Aphids, whitefly and melon thrips in various crops**

Do not apply RECRUIT 200 SC (or other Group 4A insecticides) in consecutive sprays within and between seasons. Rotate with registered insecticides from other mode of action groups.

### MIXING/APPLICATION

Prior to pouring, shake container vigorously, then add the required amount of RECRUIT 200 SC to water in the spray vat while stirring or with agitators in motion.

### EXPORT OF TREATED PRODUCE

Growers' should note that MRLs or import tolerances do not exist in all markets for edible produce treated with RECRUIT 200 SC. If you are growing edible produce for export, please check with Indigo Specialty Products Pty Ltd for the latest information on MRLs and import tolerances before using RECRUIT 200 SC.

### COMPATIBILITY

RECRUIT 200 SC is compatible with propineb, bitertanol or methamidophos. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Indigo Specialty Products Pty Ltd, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Indigo Specialty Products Pty Ltd cannot be responsible for the behaviour of such mixtures.

### PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. DO NOT graze any treated area, or cut for stock food. DO NOT feed produce or turf clippings from treated area to animals, including poultry.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways and drains with the chemical or used containers.

## **STORAGE & DISPOSAL**

Store in the closed, original containers in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

## **SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use wear cotton overalls buttoned to the neck and wrist or equivalent clothing and elbow-length chemical resistant gloves and goggles. Wash hands after use. After each day's use wash gloves and contaminated clothing.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## **SAFETY DATA SHEET**

Additional information is listed in the safety data sheet (SDS). A safety data sheet for ProForce RECRUIT 200 SC Insecticide is available from Indigo Specialty Products Pty Ltd on request. Call Customer Service on 02 9389 2455

**CONDITIONS OF SALE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by Indigo Specialty Products Pty Ltd or under abnormal conditions. Indigo Specialty Products Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.



**Indigo Specialty Products** ABN 15 631 459 660  
6/163-173 McEvoy Street, Alexandria NSW 2015  
Ph: 0402 725 887

APVMA Approval No.: 89271/126025